



**UNIVERSITI PUTRA MALAYSIA**

**THE EFFECT OF BENZO(a)PYRENE ON MALE MOUSE  
*MUS MUSCULUS***

**PARICHEHR HANACHI**

**FSAS 2002 24**

**THE EFFECT OF BENZO(a)PYRENE ON MALE MOUSE  
*MUS MUSCULUS***

**By**

**PARICHEHR HANACHI**

**Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia,  
in Fulfillment of the Requirements for the Degree of Doctor of Philosophy**

**June 2002**

## **DEDICATION**

Dedicated to my husband, Mahmood Abaie, and my daughter Somayeh for their devotion, understanding and support during all difficulties and to my parents for their true love, constant trust and principles that guided my life and to many researchers whose have not given due recognition for the many hours spent in the laboratory and friends to provide humanity with solutions to better life

Abstract of thesis presented to the Senate of Universiti Putra Malaysia in fulfilment  
of the requirement for the degree of Doctor of Philosophy

**THE EFFECTS OF BENZO(a)PYRENE ON MALE MICE  
*MUS MUSCULUS***

**By**

**PARICHEHR HANACHI**

**June 2002**

**Chairman: Assoc. Prof. Dr. Nor Aripin Shamaan**

**Faculty: Science and Environmental Studies**

A study was carried out to determine the effect of benzo(a)pyrene (BaP) on selected enzyme activities , the pathology of lung, liver and kidney and cellular aspects of the mice *Mus musculus*. A suppression or change in the activities of several enzymes in these tissues can be used as a potential technique for the diagnosis of carcinogenesis in the early stage.

The initial work involved the evaluation of lethal dose and the threshold dose required for induction of carcinogenesis in adult mice. Subsequent work involved the determination of glutathione S-transferase (GST) and glutathione peroxidase (GPx) activities over a time period during which the mice were treated with benzo(a)pyrene. Finally, induction of GST and GPx was achieved by long-term treatment with BaP. The GST was purified partially by affinity chromatography. Determination of the effect of BaP on the histology of liver, lung, kidney of mice

were also carried out. Results obtained showed that GST and GPx activities were induced by BaP in a dose-dependent manner ( 100-250 mg/kg body weight). The mice were injected with BaP at a dose of 200 mg/kg body weight once at the start of the short term study, GST activity was induced at maximum after 4 days, and after that the activity dropped to almost normal values.

The effect of BaP on GST and GPx activities in the liver, lung , kidney and blood of male mice were studied in long-term study. The mice with injected 200 mg/kg BaP once a week for 8 weeks. The GST activity was significantly increased ( $P<0.05$ ) after 2 weeks in liver, lung, kidney and blood while GPx activity was significantly increased ( $p<0.05$ ) in liver and lung with hydrogen peroxide as the substrate after 4 and 8 weeks, GPx activity did not change when cumene hydroperoxides was used as a substrate.

Histological changes in the liver and lung was observed after 2 weeks and in the kidney after 4 weeks of treatment. The kidney showed mild inflammation after 4 weeks. Liver histology of mice treated after 2 weeks showed some cells with binucleation and after 4 weeks showed degeneration and necrosis and hepatocytes were slightly enlarged. The lung cells showed severe acute inflammation after 2 weeks and after 4 weeks showed sever epitelization and the cells lost their normal shape and arrangement and the nuclei become hyperchromatic after 8 weeks.

Attempts to purify GST in the mice were carried out using GSH-sepharose affinity matrix. The cytosolic GST purified in this study resolved into three discrete molecular species of approximate molecular weight 25271 D, 23478 D and 25839 D respectively comparable to the previously designated isoforms MI, MII and MIII. The GST in the BaP treated sample exhibited electrophoretic migration on SDS-PAGE closely similar to the normal control. Purified liver GST had higher specific activity than the lung and the subunits were of comparable size. This may indicate the existence of common GST isoform in both organs. The main finding in this research related to result of IEF. It showed three activity peaks in the normal control and BaP treated samples livers with different pIs and substrate specificities. The GST activity toward ethacrynic acid in the treated mice was significantly higher than normal control indicating BaP induced the GSTMII (class Pi). In this study, laboratory trials with biochemical measurements supported by toxicity and histological studies were tested as tools for the assessment of the environmental hazard of BaP to target organisms.

Abstrak tesis dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan bagi Ijazah Doktor Falsafah

**KESAN BENZO(a)PIREN TERHADAP TIKUS JANTAN (*MUS MUSCULUS*)**

**Oleh**

**Parichehr Hanachi**

**Jun 2002**

**Pengerusi : Prof. Madya Dr. Nor Aripin Shamaan**

**Fakulti : Sains dan Pengajian Alam Sekitar**

Suatu kajian telah dijalankan mengenai kesan benzo(a)piren terhadap aktiviti beberapa enzim, patologi paru-paru, hati dan buah pinggang, dan aspek sel tikus *Mus musculus*, serta potensi memperoleh suatu teknik diagnostik untuk mengesan karsinogenesis pada peringkat awal.

Kerja selidik peringkat permulaan melibatkan penilaian dos maut dan dos ambang (threshold dose) yang diperlukan untuk mengaruh karsinogenesis bagi tikus dewasa. Kerja selidik berikutnya melibatkan penentuan/penganggaran aktiviti glutathione S-transferase dan glutathione peroksidase dalam jangka masa tikus tersebut menerima dos benzo(a)piren. Pada peringkat akhir kerja selidik, pengaruh/induksi GST dan GPx dicapai melalui pemberian dos BaP dalam jangka masa panjang. GST telah ditulen separa dengan kaedah kromatografi afiniti.. Kerja selidik tambahan juga dilakukan untuk menentukan kesan BaP terhadap histologi tisu hati, paru-paru dan buah pinggang tikus tersebut. Keputusan yang diperolehi

menunjukkan aktiviti GST dan GPx telah diaruh oleh BaP dan pengaruh bergantung kepada dos yang diberi (100-250 mg/kg berat badan). Tikus tersebut telah disuntik dengan BaP sekali pada dos 200 mg/kg berat badan pada permulaan kajian jangka pendek. Aktiviti GST diaruh ke tahap maksimum selepas 4 hari, dan selepas itu aktiviti enzim menurun ke tahap normal/lazim.

Kesan BaP terhadap GST dan GPx tisu hati buah pinggang dan darah tikus dewasa telah dikaji secara pengajian jangka panjang. Tikus tersebut telah disuntik dengan 200mg/kg BaP sekali seminggu selama 8 minggu. Aktiviti GST meningkat secara signifikan ( $P < 0.05$ ) selepas 2 minggu bagi tisu hati, paru-paru, buah pinggang dan darah. sementara aktiviti GPx meningkat secara signifikan ( $p < 0.05$ ) dalam hati dan paru-paru menggunakan  $H_2O_2$  sebagai substrat. Selepas 4 dan 8 minggu, aktiviti GPx tidak berubah apabila  $cuCOOH$  digunakan sebagai substrat.

Perubahan histologi pada hati dan paru-paru dapat diperhatikan selepas 2 minggu rawatan dan pada ginjal selepas 4 minggu rawatan. Ginjal menunjukkan berlakunya sedikit keradangan selepas 4 minggu rawatan. Selepas 2 minggu rawatan, histologi pada hati tikus menunjukkan sesetengah sel mengalami binukleasi dan selepas 4 minggu, ia menunjukkan kemerosotan manakala nekrosis dan hepatosit menjadi besar sedikit demi sedikit. Sel paru-paru pula menunjukkan berlakunya serangan radang akut selepas 2 minggu. 4 minggu selepas itu ia menunjukkan



berlakunya pematangan epitelization dan seterusnya sel mengalami kehilangan bentuk dan susunan asalnya. Nukleus pula menjadi hiperkromatik selepas 8 minggu.

Percubaan untuk menuliskan GST pada tikus telah dilakukan menggunakan matrik afiniti GSH-sepharose. GST sitosol yang ditulen dalam kerja selidik ini dapat dipisah kepada tiga spesis molekul yang berasingan, dan bersaiz 25271D, 23478 D dan 25839 D. Ia adalah dahulunya bersamaan dengan designasi MI, MII dan MIII. GST dalam sampel yang telah dirawat mempamerkan migrasi/ pergerakan elektroforesis SDS-PAG seperti kawalan normal. GST yang telah dituliskan dari hati mempunyai aktiviti spesifik yang lebih tinggi berbanding dengan paru-paru. Subunit GST dari organ tersebut merupakan saiz yang setanding. Ini menunjukkan kewujudan isoform yang biasa di dalam kedua-dua organ tersebut. Keputusan dari kaedah IEF menunjukkan tiga puncak aktiviti bagi sampel kawalan dan rawatan yang mempunyai pI berbeza dan spesifisiti substrat yang berbeza. Aktiviti GST terhadap asid ethacrynic didalam tikus telah meningkat berbanding tikus kawalan. Dalam kerja selidik ini, percubaan menggunakan pengukuran biokimia disokong dengan kajian toksiti/keracunan dan histologi telah diuji sebagai alat untuk menilai ancaman persekitaran BaP terhadap organisma sasaran.

## ACKNOWLEDGEMENTS

Words are not enough to express my sincere appreciation for the people who were involved in my training towards obtaining this degree. I would like to express my deepest appreciation to my mentor and advisor, Assoc. Prof. Dr. Nor Aripin Shamaan for his continuing encouragement and support during my Ph.D. I am very thankful for his guidance, fruitful discussions and criticisms.

I also wish to thank my committee members Assoc. Prof. Dr. Mohd Arif Syed, Assoc. Prof. Dr. Juzu Hayati Arshad and Assoc. Prof. Dr. Johari Ramli for the considerable time they invested in reading and commenting on this dissertation.

I would also like to extend my heartfelt thanks to Prof. Datin Dr. Khatijah Mohd. Yusoff and Mr. Ismail Omar for providing some of facilities, constructive suggestions and encouragement. Special thanks to Dr. Hairuszah Ithnin for all the scientific discussions and valuable comments on Histopathological studies.

I also wish to acknowledge Zarifians family and Dr. Paymon Roustaian for their friendship, enthusiastic support and companionship and Joyce, Mohamad, Dr Isam, Dr Shakinaz, Daryani, Dr Lailan Syaufina for their help and concern.

I thank the technical staff of the Department of Biochemistry and Microbiology, UPM; Ibrahim, Mohd Zamros, Onn, Husain and Khalid.

And my husband, Mahmood deserves my most heartfelt thanks and recognition for his unfailing support and belief in my ability to obtain a Ph.D., and for putting up with years of sacrifice while I was in school. Without his support and love completion of this degree would have been unlikely.

And finally, my loudest thanks to my daughter, Somayeh. Without her encouragement and uncanny and ability to keep me grounded in reality, I certainly would not have had the energy to complete this journey.



## TABLE OF CONTENTS

	<b>Page</b>
<b>ABSTRACT</b>	<b>iii</b>
<b>ABSTRAK</b>	<b>vi</b>
<b>ACKNOWLEDGEMENTS</b>	<b>ix</b>
<b>APPROVAL</b>	<b>xi</b>
<b>DECLARATION</b>	<b>xiii</b>
<b>LIST OF TABLES</b>	<b>xvii</b>
<b>LIST OF FIGURES</b>	<b>xix</b>
<b>LIST OF PLATES</b>	<b>xxii</b>
<b>LIST OF ABBREVIATIONS</b>	<b>xxiv</b>
 <b>CHAPTER</b>	
<b>I      INTRODUCTION</b>	<b>1</b>
 <b>II     LITERATURE REVIEW</b>	<b>5</b>
Cancer in General	5
Multistage Carcinogenesis	6
Chemical Carcinogen	7
Classes and Metabolism of Chemical Carcinogenesis	8
Polycyclic Aromatic Hydrocarbons	9
Benzo(a)pyrene	10
Mechanism of Pyrolytic Formation of BaP	11
Absorption and Distribution	13
Mechanism of Action	14
Metabolism	16
Biotransformation	16
Marker Enzyme	18
Glutathione S-Transferase (GST)	19
Importance of GST Pi as Marker of Carcinogenesis	22
Glutathione Peroxidase (GPx)	24
Glutathione Reductase (GRx)	25
γ-glutamyl transaminase (GGT)	26
The Glutathione Redox Cycle	27
Target Organ Toxicity	29
Liver	29
Liver Cancer	30
Histological Alterations in Hepatocarcinogenesis	32



	Liver Necrosis	33
	Lung	34
	Lung Cancer	36
	Kidney	37
	Kidney Damage	38
	The Proteins of Normal and Neoplastic Tissue	39
	Methods of Purification and Characterisation of GST	42
<b>III</b>	<b>GENERAL MATERIALS AND METHODS</b>	45
	Experimental Design	45
	Chemicals	45
	Animals	46
	Intraperitoneal Injection	46
	Preparation of Cytosol and Microsome	47
	Glutathione S-Transferase Assay	48
	Glutathione Peroxidase Assay	49
	Protein Determination	50
	Purification of GST	52
	Ammonium Sulphate Precipitation	53
	Glutathione Agarose Affinity Chromatography	53
	Electrophoresis	54
	SDS PAGE	54
	NATIVE PAGE	56
	Isoelectric-Focusing	57
	Two Dimensional Electrophoresis	59
	Isoelectric Focusing ( Aqueous Media)	60
	Histology Studies	61
	Statistic Analysis	61
<b>IV</b>	<b>EFFECT OF DIFFERENT DOSE OF BENZO(a) PYRENE ON GLUTATHIONE TRANSFERASE AND GLUTATHIONE PEROXIDASE ACTIVITIES IN MICE LIVER</b>	
	Introduction	62
	Material and Method	63
	Result	65
	Discussion	71
<b>V</b>	<b>EFFECT OF BaP ON GLUTATHIONE TRANSFERASE AND GLUTATHIONE PEROXIDASE ACTIVITIES IN MICE LIVER IN A SHORT TERM STUDY</b>	
	Introduction	74
	Material and Methods	76

	Result	79
	Discussion	86
<b>VI</b>	<b>PURIFICATION OF GLUTATHIONE TRANSFERASE IN CONTROL AND BENZO (a)PYRENE TREATED FOR TWO WEEKS</b>	
	Introduction	90
	Material and Method	91
	Result	94
	Discussion	108
<b>VII</b>	<b>EFFECT OF BaP ON MICE LIVER, LUNG, KIDNEY AND BLOOD IN A LONG TERM STUDY</b>	
	Introduction	112
	Material and Method	113
	Result	117
	Discussion	132
<b>VIII</b>	<b>PURIFICATION AND CHARACTERISATION OF GST IN CONTROL AND MICE TREATED WITH BaP FOR 2 MONTHS</b>	
	Introduction	137
	Material and Method	139
	Result	142
	Discussion	155
<b>IX</b>	<b>GENERAL DISCUSSION AND CONCLUSION</b>	160
	<b>BIBLIOGRAPHY</b>	174
	<b>APPENDICES</b>	
	1- Reagent for Glutathione S-Transferase	194
	2- Reagent for Glutathione Peroxidase	195
	3- Reagent for Protein Determination	196
	4- Preparation Affinity Chromatography Solution	197
	5- Tissue Processing by an Automatic Tissue Processor	199
	6- Preparation SDS-PAGE Solution	200
	7- Preparation IEF Solution	202
	<b>BIODATA OF THE AUTHOR</b>	204

## LIST OF TABLE

Table	Page
1 Causes of cancer	7
2 Rat cytosolic GST subunits	20
3 Histological classification of brochogenic carcinoma	36
4 Composition of running and stacking SDS-PAGE gels	55
5 Effect of different BaP doses on body weight in control and treated mice after two weeks	67
6 The specific activity of GST and GPx after 2 weeks in the liver of mice	68
7 The specific activity of GST and GPx at 200 mg/kg BaP over time in the mice liver.	80
8 Purification of GST at 200 mg/kg BaP exposure in liver of control and treated mice by affinity chromatography.	98
9 Purification of GST at 200 mg/kg BaP exposure in lung of control and treated mice by affinity chromatography	99
10 The analysis of pI values and type of activity of control liver and Lung GST at 200 mg/kg BaP exposure in short term study in mice.	107
11 The effect of BaP on body weight and liver, kidney, lung weight of mice after 8 weeks.	118
12 The specific activity of GST and GPx at 200 mg/kg BaP over time in the plasma, liver, kidney, lung of mice.	119
13 Purification by Affinity Chromatography of liver GST from control and BaP treated mice (long term study).	143
14 Specific activity with selected substrates of three major peak (affinity chromatography) GST in control and mouse treated with 200 mg/kg BaP.	147



15	Some of the compounds commonly used as substrate in the spectrophotometric method assay for GST activity.	154
16	Tissue processing by automatic tissue processor.	199



## LIST OF FIGURE

Figure	Page
1 Multistage of cancer.	5
2 Typical polycyclic aromatic hydrocarbons.	10
3 Mechanism of pyrolytic formation of BaP from organic materials.	12
4 The detoxification and metabolic activation of BaP.	15
5 Integration of phase I and phase II enzyme in carcinogens.	17
6 GST catalyse cumene hydroperoxide.	21
7 Detoxification of BaP with GST to form water soluble metabolite.	23
8 Diagram of hepatic cell arrangement in the liver.	29
9 Basic organisation of the respiratory tract.	35
10 Summary diagram of methods used to measure the specific activity of marker enzyme.	47
11 Summary diagram of methods used for purification of GST in mice liver.	53
12 Summary diagram of the methods used to prepare the samples for measuring various enzyme	64
13 Comparison of percentages changes of GST and GPx activity in male mice liver two weeks after treatment with different doses of BaP.	69
14 Effect of BaP on enzyme specific activity of GST and GPx in mice liver two weeks after treatment with different doses of BaP.	70
15 Summary diagram of the methods used to prepare the samples for measuring GST and GPx activities.	78
16 Comparison of percentage changes of GST and GPx activities in mice liver 2 weeks after treatment with 200 mg/kg BaP.	81

17	A short term study on the effect of 200 mg/kg BaP on specific Activity of liver GST.	82
18	A short term study on the effect of 200 mg/kg BaP on specific activity of liver GPx (Sub= H <sub>2</sub> O <sub>2</sub> ) .	83
19	A short term study on the effect of 200 mg/kg BaP on specific activity of liver GPx ( Sub=cuOOH ).	85
20	Summary diagram of the purification of GST in mice liver and lung.	93
21	Total protein and GST activity in supernatant of ammonium sulphate precipitation.	95
22	Total protein and GST activity in pellet of ammonium sulphate precipitation.	95
23	Profile for the purification of GST in mice liver using GSH-affinity chromatography in short term study.	100
24	Profile for the purification of GST in mice lung using GSH-affinity chromatography in short term study.	100
25	Summary diagram of the methods used to prepare the samples for measuring GST and GPx activities and histology .	115
26	A long term study on the effect of BaP on the GST activity in liver, lung, kidney and blood.	120
27	A long term study on the effect of BaP on the GPx ( Sub=H <sub>2</sub> O <sub>2</sub> ) activity in liver, lung, kidney and blood.	122
28	A long term study on the effect of BaP on the GPx ( Sub= cuOOH) activity in liver, lung, kidney and blood.	123
29	Summary diagram of the methods used to prepare the samples for purification and characterisation of GST.	140
30	Profile for the purification of GST in control mice liver using GSH- affinity chromatography.	144

31	Profile for the purification of GST in treated mice liver using GSH affinity chromatography.	144
32	Elution profile of Isoelectric focusing column loaded with partially purified GST from affinity chromatography in control mice liver.	148
33	Elution profile of Isoelectric focusing column loaded with partially purified GST enzyme sample from affinity chromatography in treated mice liver.	148
34	GST specific activity with selected substrate of three major Peaks from IEF in control mice liver.	152
35	GST specific activity with selected substrate of three major peaks from IEF in treated ( 200 mg/kg BaP) mice liver in two months.	152
36	Comparison of percentage changes of GST activities with selected substrate of three major peak from IEF in the liver mice induced by 200 mg/kg BaP in two months.	153

## LIST OF PLATES

Plates		Page
1	SDS- PAGE of control and treated mouse liver and lung with 200 mg/kg BaP in two weeks.	102
2	Native PAGE of control and treated mouse liver with 200 mg/kg BaP. Liver GST preparation over 2 weeks.	102
3	Analysis of 100,000g homogenate of control liver with two dimensional electrophoresis after two weeks.	103
4	Isoelectric Focusing PAGE using 10 $\mu$ l IEF standard with different running time.	104
5	Analysis of control and treated liver and lung sample in IEF-PAGE after two weeks.	106
6	A photomicrograph of liver from control mice .	126
7	A photomicrograph of liver which received dose 200 mg/kg BaP every week over period of 2 weeks.	126
8	A photomicrograph of liver received dose 200 mg/kg BaP every week over period of 4 weeks.	127
9	A photomicrograph of liver of mice which received a dose 200 mg/kg BaP every week over a period of 8 weeks.	127
10	Kidney section from control mice .	128
11	A photomicrograph of the kidney of mice which received 200 mg/kg BaP every week over period of 2 weeks.	128
12	A photomicrograph of kidney of mice which received a dose of 200 mg/kg BaP every weeks over a period of 4 weeks.	129
13	A photomicrograph of kidney of mice which received 200 mg/kg BaP every week over a period of 8 weeks.	129

14	Lung section from control mice .	130
15	A photomicrograph of the lung of a mice which received 200 mg/kg BaP per week over period of 2 weeks.	130
16	A photomicrograph of lung mice which received 200 mg/kg BaP per week over a period of 4 weeks .	131
17	A photomicrograph of lung of mice which received 200mg/kg BaP per weeks over period of 8 weeks .	131
18	SDS PAGE of GST at 200 mg/kg BaP exposure in treated and control mice liver after two months.	146
19	SDS PAGE gel profiles of control mouse liver cytosolic GST purified by affinity binding GSH column after two months.	149
20	SDS PAGE gel profiles of treated (200 mg/kg BaP) mouse liver cytosolic GST purified by affinity binding to the GSH column after two months.	149

## LIST OF ABBREVIATIONS

AAF	2-Acethylaminofluorene
APS	Ammonium persulphate
a.a	Amino Acid
BSA	Bovine Serum Albumin
BaP	Benzo(a)pyrene
BPB	Bromophenol blue
Bp	Boiling point
C	Control
CV	Coefficient of variation
CDNB	1-chloro-2,4-dinitrobenzene
cuOOH	Cumen hydroperoxides
2DE	2 Dimensional Electrophoresis
DCNB	1,2-Dichloro-4-nitrobenzene
DEN	Diethylnitrosamine
DNA	Deoxyribonucleic acid
DEAE	Diethylaminoethyl
e.u.	Enzyme per unit
EA	Ethacrynic acid
EDTA	Ethylenediaminetetra acetic acid
GST	Glutathion S-transfrase
GPx	Glutathion peroxidase
GSH	Glutathione (reduced form)
GSSG	Glutathione disulphate (oxidised)
g	Gram
HCl	Hydrogen chloride
IEF	soelectric focusing
K <sub>2</sub> HPO <sub>4</sub>	Potassium hydrogen phosphate
KH <sub>2</sub> PO <sub>4</sub>	Potassium dihydrogen phosphate
KCl	Potassium chloride
L	liter
mA	Miliampere
mg	Milligram
mL	Millilitre
mM	Millimolar
mw	Molecular weight
mp	Melting point
M	Molar
ME	Mercaptoethanol
NADH	Reduced nicotinamide adenine dinucleotide

