



**UNIVERSITI PUTRA MALAYSIA**

**REVIEW ON SYSTEMATIC OF CALANOID COPEPOD (CRUSTACEA)  
FROM SEAGRASS AND CORAL AREAS OF THE  
EAST COAST OF THE MALAY PENINSULA**

**MOHD AFANDY BIN BABA @ ABDUL KADIR**

**FSAS 2002 39**

**REVIEW ON SYSTEMATIC OF CALANOID COPEPOD (CRUSTACEA)  
FROM SEAGRASS AND CORAL AREAS OF THE  
EAST COAST OF THE MALAY PENINSULA**

**By**

**MOHD AFANDY BIN BABA @ ABDUL KADIR**

**Thesis submitted to the School of Graduates Studies, Universiti Putra Malaysia,  
in Fulfillment of The Requirements for the Degree of Master of Science**

**April 2002**



**DEDICATION**

**MY PARENT (& IN-LAW)  
ABAH, MAK AND MEK**

**MY BELOVED WIFE,  
ZAHARAH CHE MAHMOOD.**

**MY LITTLE LOVELY SONS AND DAUGHTER,  
MUHAMMAD NASYRUL ADLY  
MUHAMMAD AIMAN SYAHMY  
NURUL IFFAH  
EMRAN NUR HAKIM**

**MY SISTERS AND BROTHERS (& IN-LAWS)  
ALL MY TEACHERS,  
OR THOSE WHO GIVE ME KNOWLEDGES,  
ALL MY FRIENDS**

**Thank you for all your prayers and endless support**

**JAZAKUMULLAHU KHAIRAN KATHIIRA**

Abstract of thesis presented to the Senate of Universiti Putra Malaysia in fulfillment of the requirements for the degree of Master of Science.

**REVIEW ON SYSTEMATIC OF CALANOID COPEPOD (CRUSTACEA)  
FROM SEAGRASS AND CORAL AREAS OF THE  
EAST COAST OF MALAY PENINSULA**

By

**MOHD AFANDY BIN BABA @ ABDUL KADIR**

**April 2002**

**Chairman : Drs. Idris Abdul Ghani**

**Faculty : Science and Environmental Studies**

A study has been carried out on the systematic of calanoid copepod from samples taken off sea grass and coral area in East Coast of Peninsular Malaysia. The material examined was taken from various samples collected since 1990 until 1997 in Pulau Sibü, Pulau Besar, Pulau Redang and Pulau Bidong. Thirty-four species representing 16 genera and 12 families were described and illustrated. Six species are new record for Malaysian coastal water. They are *Subeucalanus crassus*, *S. subcrassus*, *S. pileatus*, *Labidocera kroeyeri*, *Centropages yamadai* and *Bestiolina similis*. *Subeucalanus* is a new record for genus. This study also found that the species *Centropages yamadai* is also new to Indo Pacific region.

This study also reviewed the systematic and distribution of calanoid copepod in Malaysian waters based on published and unpublished literatures. Misidentification and misuse of species names in the literature have been corrected where possible.

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk ijazah Master Sains

**KAJIAN SEMULA KE ATAS SISTEMATIK BAGI KOPEPOD CALANOID  
(CRUSTACEA) DI KAWASAN RUMPUT LAUT DAN TERUMBU KARANG  
PANTAI TIMUR SEMENANJUNG MALAYSIA.**

Oleh

**MOHD AFANDY BIN BABA @ ABDUL KADIR**

**April 2002**

**Pengerusi : Drs. Idris Abdul Ghani**

**Fakulti : Sains dan Pengajian Alam Sekitar**

Sistematik calanoid kopepod di ekosistem rumput laut dan batu karang di perairan pantai timur Semenanjung Malaysia telah diteliti. Sampel kajian adalah hasil pengumpulan beberapa siri penyampelan yang dilakukan dari tahun 1990 hingga 1997 di Pulau Sibul, Pulau Besar, Pulau Redang dan Pulau Bidong. Sebanyak tiga puluh empat spesis mewakili 16 genera dan 12 famili yang telah dikenalpasti dan dilukis. Enam spesis merupakan rekod baru bagi perairan Malaysia. Spesis tersebut ialah *Subeucalanus crassus*, *S. subcrassus*, *S. pileatus*, *Labidocera kroeyeri*, *Centropages yamadai* dan *Bestiolina similis*. Kajian ini juga mendapati spesis *Centropages yamadai* adalah rekod baru kepada perairan Indo-Pacific.

Kajian ini juga mengulas kembali tentang sistematik and taburan kopepod calanoid di perairan Malaysia berdasarkan bahan yang telah diterbitkan atau tidak diterbitkan. Ketidaktepatan pengecaman dan kesilapan pada nama spesis telah dibetulkan sekiranya perlu.

## ACKNOWLEDGEMENTS

In the name of Allah, the Most Gracious and the Most Merciful. Praise is to Allah, the Cherisher and Sustainer of the worlds, for giving me the toughness to complete my thesis.

The greatest thankful and appreciation to my supervisor and supervisory committee, Drs Hj. Idris Abdul Ghani, Dr Hishamuddin Omar and Dr Misri Kusnan. I also would like to thank Assoc. Professor Dr Ahmad Ismail and Dr Abdul Rahim Ismail from Biology Department, UPM for their kindness guidance; Dr. Bradford from New Zealand Institute of Water Association (NIWA), Dr. Razouls from Institute' de Oceanographique, Canada and Dr Susumu Ohtsuka from University of Tokyo, Japan for their greatest cooperation's and guidance on copepodology. Thanks are also due to all the members of Aquatic Laboratory and all my friends in Biology Department, UPM.

Lastly, I would like to appreciate the understanding of my wife who had sacrificed and encouraged me to be excellent in many aspects.

## TABLE OF CONTENTS

	Page
DEDICATION	ii
ABSTRACT	iii
ABSTRAK	iv
ACKNOWLEDGEMENTS	v
APPROVAL LETTER	vi
DECLARATION	viii
LIST OF TABLES	xi
LIST OF FIGURES	xii
LIST OF ABBREVIATIONS	xv
 <b>CHAPTER</b>	
<b>1. INTRODUCTION</b> .....	1.1
1.1 Importance of Study .....	1.3
 <b>2. LITERATURE REVIEW</b> .....	2.1
2.1 Seagrass and Coral Reef Habitat .....	2.2
2.2 Classification of Calanoid Copepod .....	2.4
2.2 Taxonomic Composition and Distribution.....	2.7
2.4 Studies in Malaysian Coastal Water.....	2.10
 <b>3. MATERIALS AND METHODS</b> .....	3.1
3.1 Study Area and Sampling Program .....	3.1
3.2 Laboratory Works .....	3.3
3.3 Species Identification and Drawings.....	3.5
3.4 Body Length Measurement.....	3.6
 <b>4. RESULTS</b> .....	4.1
4.1 Systematic Species Composition .....	4.1
4.2 Provisional Identification Keys For Malaysian Calanoid Copepod .....	4.3
4.2.1 Key to the Family of Calanoida .....	4.3
4.2.2 Family CALANIDAE Giesbrecht, 1888 .....	4.4
4.2.3 Family PARACALANIDAE Giesbrecht, 1892 .....	4.9
4.2.4 Family CLAUSOCALANIDAE Giesbrecht, 1892 .....	4.19
4.2.5 Family EUCHAETIDAE Giesbrecht, 1892 .....	4.22
4.2.6 Family EUCALANIDAE Giesbrecht, 1892 .....	4.24
4.2.7 Family CALOCALANIDAE Bernard, 1958 .....	4.30
4.2.8 Family: ACARTIDAE Dana, 1846 .....	4.34
4.2.9 Family CENTROPAGIIDAE Giesbrecht, 1892 .....	4.39
4.2.10 Family PONTELLIDAE Dana, 1853 .....	4.47
4.2.11 Family TEMORIDAE Giesbrecht, 1892 .....	4.62
4.2.12 Family: TORTANIDAE Sars, 1902.....	4.68
4.2.13 Family PSEUDODIAPTOMIDAE G.O.Sars, 1902.....	4.72
4.3 Biodiversity of Calanoid Copepod in Malaysian Marine Waters.....	4.74
4.3.1 Family CALANIDAE Giesbrecht, 1888.....	4.77
4.3.2 Family PARACALANIDAE Giesbrecht, 1892.....	4.80
4.3.3 Family CLAUSOCALANIDAE Giesbrecht, 1892.....	4.81
	ix

4.3.4 Family EUCHAETIDAE Giesbrecht, 1892.....	4.82
4.3.5 Family EUCALANIDAE Giesbrecht, 1892.....	4.82
4.3.6 Family CALOCALANIDAE Bernard, 1958.....	4.82
4.3.7 Family: ACARTIDAE Dana, 1846.....	4.83
4.3.8 Family CENTROPAGIIDAE Giesbrecht, 1892.....	4.84
4.3.9 Family PONTELLIDAE Dana, 1853 .....	4.85
4.3.10 Family TEMORIDAE Giesbrecht, 1892 .....	4.86
4.3.11 Family: TORTANIDAE Sars, 1902 .....	4.86
4.3.12 Family PSEUDODIAPTOMIDAE G.O.Sars, 1902 .....	4.87
4.3.13 New Recorded Species .....	4.87
<b>5.0 CONCLUSION</b> .....	<b>5.1</b>
<b>BIBLIOGRAPHY</b>	
<b>VITAE</b>	



## LIST OF TABLES

Table		Page
3.1	List of Sampling Stations	3.2
4.1	Species occurrence of Calanoid Copepod in Study Areas.	4.75
4.2	Comparison on the seta formula of dominant species in all study areas	4.78
4.3	Morphological Comparison on the appendages between <i>Subeucalamus</i> and <i>Eucalamus</i>	4.89
4.4	Comparison of species occurrence of calanoid copepod distribution within Malaysian water.	4.91

## LIST OF FIGURES

Figure		Page
1.1	Copepod habitats: representation of the primary habitat of each of the ten copepod orders. A. Platycopioidea. B. Misophrioida. C. Harpacticoida. D. Calanoida. E. mormonilloidea. F. Cyclopoida. G. Monstrilloidea. H. Poecilostomatoida. I. Siphonostomatoida. J. Gelyelloidea. [A-C, benthic; D-G, planktonic; H and I, associated; J, groundwater](Huys and Boxshall, 1991)	1.2
2.1	General Morphology of Calanoid Copepod (after Giesbrecht, 1893)	2.6
3.4	Seta formula system (after Sewell, 1947)	3.6
4.1	<i>Canthocalanus pauper</i> Giesbrecht, 1888: Female :- A – dorsal view, B – lateral view, C – leg 1, D – leg 5; Male, E – dorsal view, F – lateral view, G – leg 1, H – leg 5.	4.6
4.2	<i>Undimula vulgaris</i> Dana, 1849: Female:- A – dorsal view, B – lateral view, C – Th5-lateral view, D – Th5- lateral view (bifurcate), E – leg2, F – leg5; Male:- G – dorsal view, H – lateral view, I - leg5.	4.8
4.3	<i>Acrocalanus gibber</i> Giesbrecht, 1888: Female:- A - Dorsal view, B – lateral view, C – leg 4; Male:- D – lateral view, E - leg 5.	4.11
4.4	<i>Acrocalanus gracilis</i> Giesbrecht, 1888: Female:- A – Dorsal view, B – leg 4 (from Bradford-Grieve, 1994)	4.13
4.5	<i>Acrocalanus longicornis</i> Giesbrecht, 1888: Female:- A – Dorsal view, B – lateral view, C – leg 4. (A and C from Bradford-Grieve, 1994)	4.14
4.6	<i>Bestiolina similis</i> Sewell, 1914: Female:- A – lateral view, B – leg 4; Male:- C – lateral view of posterior body, D – leg 5 (from Bradrord-Grieve, 1994).	4.15
4.7	<i>Paracalanus aculeatus</i> Giesbrecht, 1888 (from Bradford-Grieve, 1994): Female:- A - lateral view, B – dorsal view, C – leg 4, D – leg 5; Male:- E –dorsal view, F - leg 5.	4.18
4.8	<i>Paracalanus parvus</i> Claus, 1863: Female:- A – dorsal view, B – lateral view, C – leg 4.	4.20
4.9	<i>Clausocalanus furcatus</i> Brady, 1883: Female :-A – dorsal view, B – lateral view, C – caudal rami, D – leg 5.	4.21

- 4.10 *Euchaeta concinna* Dana, 1849: Female:- A – dorsal view, B – leg 2.; Male:- C – dorsal view, D – leg 5, E – exopod of leg 3. 4.23
- 4.11 *Subeucalanus crassus* Geletin, 1976: Female:- A - dorsal view, B - Urosome (dorsal view); C – mandible, D - part of maxilla 1. 4.25
- 4.12 *Subeucalanus subcrassus* Geletin, 1976: Female:- A – dorsal view, B – urosome (dorsal view); C – mandible, D – part of maxilla 1. 4.27
- 4.13 *Subeucalanus crassus* Geletin, 1976: Female :- A - dorsal view, B – mandible, C - part of maxilla 1. 4.29
- 4.14 *Calocalanus pavo* Dana, 1849:- Female:- A – dorsal view, B – dorsal view (from Mori, (1937) - scale not provided), C – leg 4, D – leg 5. 4.31
- 4.15 *Calocalanus pavoninus*, Farran, 1936: Female:- A – dorsal view, B – dorsal view, C – leg 5, D – leg 4. (from Bradford-Grieve, 1994). 4.33
- 4.16 *Acartia erythraea* Giesbrecht 1889. Female :- A – dorsal view, B – lateral view, C – urosome, D- leg 5. Male :- E – dorsal view, F – urosome, G – leg 5. 4.35
- 4.17 *Acartia spinicauda* Giesbrecht, 1889. Female:- A – dorsal view, B – Urosome, D – leg 5. Male:- A – dorsal view, B – Urosome, C – leg 5. 4.38
- 4.18 *Centropages furcatus* Dana, 1849: Female:- A – dorsal view, B – lateral view, C – leg 5; Male:- D – dorsal view, E – leg 5. 4.40
- 4.19 *Centropages longicornis* Mori 1932: Female:- A – dorsal view, B – leg 5; Male – C – dorsal view, D – right antennule, E – leg 5. 4.42
- 4.20 *Centropages orsinii* Giesbrecht, 1889: Female:- A – dorsal view, B – lateral view, C – leg 5; Male:- D – dorsal view, D – leg 5. 4.44
- 4.21 *Centropages yamadai* Mori, 1934: Male :- A dorsal view, B – right leg 4, C – left leg 4, D – leg 5.. 4.46
- 4.22 *Calanopia elliptica* Dana 1849: Female:- A – Dorsal view, B – leg 5; Male:- C – dorsal view, D – leg 5. 4.48
- 4.23 *Calanopia thompsoni*, A. Scott, 1909: Female:- A - Dorsal view, B – lateral view, C – leg 4. 4.51
- 4.24 *Labidocera acuta* Dana, 1849 (from Bradford-Grieve, 1999): Female:- A – Dorsal view, B –leg 5; Male:- C –dorsal view, D – leg 5. 4.52

4.25	<i>Labidocera bengalensis</i> Krishnaswamy, 1952: Female:- A - dorsal view, B – leg 5; Male (from Sillas & Pillai, 1973):- C – dorsal view, D - leg 5.	4.55
4.26	<i>Labidocera minuta</i> Giesbrecht, 1889: Female:- A – Dorsal view, B – urosome, C - leg5 (from Sillas & Pillai, 1973).	4.57
4.27	<i>Labidocera pavo</i> Giesbrecht, 1889: Female:- A - Dorsal view, B – lateral view, C – leg 5.	4.59
4.28	<i>Labidocera kroeyeri</i> Thompson & Scott, 1903: Female:- A - Dorsal view, B – lateral view, C – Uro, D - lateral of posterior body, E – leg 5	4.61
4.29	<i>Temora discaudata</i> Giesbrecht, 1889: Female:- A - Dorsal view, B – lateral view, C – leg 5; Male:- D –dorsal view, E - leg 5.	4.64
4.30	<i>Temora stylifera</i> Dana, 1849: Female:- A – Dorsal view, B – Uro, C – leg 5.	4.65
4.31	<i>Temora turbinata</i> Dana, 1849: Female:- A – Dorsal view, B – leg 5; Male:- C – dorsal view, D - leg 5.	4.67
4.32	<i>Tortanus bowmani</i> Othman, 1987: Female:- A – Dorsal view, B – lateral view, C – Urosome.	4.69
4.33	<i>Tortanus forcipatus</i> Brady, 1883: Female:- A – dorso-lateral view, B – leg 5; Male:- C – dorsal view & leg 5, D – antennule. <i>Tortanus gracilis</i> Brady, 1883: Female:- A - Dorsal view, B – leg 5; Male:- D – lateral view, E – leg.	4.71
4.34	<i>Pseudodiaptomus</i> sp. : Female:- A - Dorsal view, B – lateral view, C – leg 4; Male:- D – lateral view, E - leg 5.	4.73
4.35	Appendage of <i>Eucalamus</i> ( <i>Eucalamus hyalinus</i> , Claus 1866 – from Bradford (1994). Female:- A – Uro (dorsal view), B – Mandible, C – Maxilla of <i>E. elongatus</i> (from Mori, 1937)	4.90

## LIST OF ABBREVIATIONS

A1	- Antennule/first antenna
A2	- Antenna
CR	- caudal rami
Th	- thoracic segment.
Th1, Th2, ..	- thoracic segment 1, thoracic segment 2, ....
Ab	- Abdomen
Uro	- urosome
Uro1, Uro2...	- urosome segment 1, segment 2.....
Mdb	- mandibula
M	- maxilla
M1	- maxillule
H'	- diversity index
s.d.	- standard deviation


# CHAPTER 1

## INTRODUCTION

The Malay Peninsula surrounded by the Malacca Straits in the west coast and South China Sea in the east coast, which are highly abundant with unexplored fauna, that part of the complex planktonic life forms. The taxonomical study of marine copepods had been developed since early of the 18<sup>th</sup> decade. In Malaysia study of those copepod fauna has not been well updated. The taxonomy of calanoid copepod in Malaysia has to be revised prior to other aspects of study. This study is focused on the calanoid copepods, the most dominant fauna of planktonic crustacean.

Calanoida is the biggest among ten order of the subclass Copepoda, Class Crustacea, phylum Arthropoda. They are also being the most diverse group of in term of species especially in marine, which composed over a thousand species of all over the world (Sorokin, 1993). They occupied several types of marine habitats as showed in the schematic representation of the primary habitat of each of the ten copepod orders as in Figure 1.1.

Some species of calanoid may serve as food for livestock and domestic fowls. *Calanus finmarchicus* and *Calanus plumchrus* are the most dominant species of calanoid copepod in the North Atlantic and North Pacific Ocean respectively. They are caught and utilized directly as crustacean paste for fish consumption in aquaculture. At the same time, *Calanus plumchrus* has been utilized directly for many years in Hokkaido, Japan. These marine planktonic copepods are of high nutrient value. For example calculated as dried weight, Calanus contains 10 to 40%



1.2 Figure 1.1: Copepod habitats: representation of the primary habitat of each of the ten order of copepod. A. Platycopioida. B. Misophrioida. C. Harpacticoida. D. Calanoida. E. Mormonilloida. F. Cyclopoida. G. Monstrilloida. H. Poecilostomatoida. I. Siphonostomatoida. J. Gelvelloida. [A-C, benthic; D-G, planktonic; H and I, associated; J, groundwater](Huys and Boxshall, 1991)



protein, 12 to - 47% lipid, 3% chitin and 3.6% ash. In the protein about 20 kinds of important amino acids are found (Zheng, 1989). Consequently, copepods used as food for livestock are of high economic significance.

A large number of adult fish, particularly tropical reef fish, make evasive prey such as calanoids an important part of their diet (Emery, 1973; Hobson, 1974; Hobson & Chess 1976 (in Milton *et al.*, 1990)). These reef zooplanktivores, sometimes termed water column foragers (Davies & Birdsong, 1973 (in Sorokin, 1993) of plankton pickers (Collete & Talbot, 1972) are diurnal opportunistic zooplanktivores which feed on the plankton found near coral reefs, often forming dense, multispecies aggregations of fish.

Most evidence of reef fish feeding on calanoids comes from gut content analysis of wild caught fish (e.g. Emery, 1973; Hobson & Chess, 1976 (in Milton *et al.*, 1990)). While this type of data describes the principal components of diet of zooplanktivorous fish, it does not address an equally important question; by what mechanism can some zooplanktivores feed regularly on highly evasive prey such as calanoid copepods?

### **1.1 Importance of Study**

The study of copepod had been started since the early of 18<sup>th</sup> decade but the information on copepods from Malaysian waters still rare. It has to be established and upgraded prior to any developments in all types the coastal areas due to rapid developments towards the year 2020. Subsequently, datas of copepods from coral



reef and seagrass ecosystems around Malaysian water are need to investigate. It is essential and prior for scientific research and also to wider the knowledge on the zooplankton in Malaysia waters.

Pertinent literature of the systematic of calanoid copepods of Malaysian coastal waters is not been done extensively. Previously, species of copepods from Malaysian waters have been recorded in some general studies on the distribution and ecology. Started with pioneer study by Sewell in 1933, it is then followed by Wickstead (1961); Seng, (1984); Sarepah (1987); Idris *et al.*, (1993); Zalina, (1993) and Idris *et al.*, (1997a,1997b, 1999). All those studies were mostly discussed on the distribution of calanoid copepods as a major part of marine zooplankton.

Studies on the taxonomy of Malaysian pelagic copepods are extremely rare. The only studies have been carried out by Arvin (1977); Zuraini (1986); Othman (1986, 1987) and Zaleha, (1996). The knowledge of this fauna (copepods) in Malaysian marine water region should be started with a detail study on their taxonomical description. Taxonomical studies should be the first consideration on any studies of living fauna.

More studies are needed to determine the species composition, life cycles, feeding habit, diurnal movements, lunar cycles and seasonal cycles of this unique group of plankton (Porter & Porter 1977). Obviously, many more reefs will have to be sampled before zoogeographical trends can be seen. Most 'reefs' plankton studies to date have not directly addresses to this group. According to Sorokin (1993), it must however, be emphasised that an accurate specific diagnosis is a matter for a specialist and will usually involve micro-dissection and examination of the appendages.

According to Wolfgang (1986), species differences are often minute and superficial examination without careful study of original literature only allows gross identification.

The objectives of this study are as follows:-

1. To update the record of calanoid copepod biodiversity in Malaysian coastal water especially at sea grass and coral area.
2. To establish the taxonomical characteristic descriptions and illustrations calanoid copepods in coral reefs and sea grass ecosystems within Malaysian marine waters.
3. To review the systematic and distribution of calanoid copepods in Malaysian waters.

## **CHAPTER 2**

### **LITERATURE REVIEW**

#### **2.1 Introduction to the Calanoid Copepod.**

As the most diverse group amongst copepod, the order of Calanoida is also known as a dominant zooplankton and constitutes of a considerable part of the secondary production. They are abundant in fresh water and inland saline habitats in British waters or, for that matter in most other sea areas (Newell & Newell, 1977).

Most of Calanoid copepods are generally to be pelagic, living in the water column away from the influence of the seabed. Some live in the near-bottom community and a few are more or less benthic (Huys & Boxshall, 1991), which known as hypermetric or demersal Calanoid (Ohtsuka, 1992). They occur at all depths in the water column, from surface to the abyssal trenches.

Zooplankton is a very important biota in the ocean not only do they serve as primarily uses; they are also food for economic fishes. Zooplankton is also sensitive to any environmental changes thus it can be a good indicator to access the changes in marine ecosystem.

The environment and plankton community together forms comprehensive ecosystem as in any aquatic ecosystem. These interactions give rise to many adaptations by the organism to a planktonic existence (Wickstead, 1976)



In general copepods are the major members of the marine zooplankton community, constituting the food item of fishes. Hence, copepod occupies an important position in the food web and energy flows of the marine ecosystem environments.

Many crustaceans are phytoplankton feeder and pelagic economic fishes and whales then feed them upon. Hence, planktonic crustaceans play an important role in the food cycle of the sea. Copepoda, Ostracoda and Malacostraca are three of the important subclasses of this phylum (Zheng, 1989). Among the important group in subclass copepoda are the order calanoid and cyclopoid and others.

On looking at the potential at the copepod being a replacement to *Artemia*, the fish Fisheries Department of Malaysia is conducting research of monoculture of copepod in big tanks and brackish ponds (Azhar, 1998). This copepod will then supplied to the hatcheries. Copepod such as *Acartia*, *Paracalanus*, *Pseudodiaptomus* and *Oithona* are good prey for fish larvae.

## **2.1 Seagrass and Coral Reef Habitat**

The Indo-West Pacific Basin is concentrated the greatest number of species of marine life (Ekman, 1991). Being a part of the faunistic region of the Indo-West Pacific, Malaysia blessed with a rich marine environment and ecosystems such as coral reefs, sea grasses, seaweeds, mangroves and various kinds of beaches. They are known to associate with various kinds of fauna especially in the aquatic environment. One of them is the zooplankton, which occurred, in a bulk number in term of species as well as the individuals.

Sea grass and coral reefs are known for the diversity and abundance of their fish and invertebrate faunas. Sea grass beds are recognized as important nursery ground for many important commercial and forage organisms as well as some of the reef species. About 12% of the total fish catches in the world are produced by coral reefs and sea grass ecosystem (Fortes, 1990). These ecosystems provide food and shelter to avoid strong current, predator and sunlight to various of marine faunas.

Sea grass inhabitants of tropical and temperate waters, form important part of coastal ecosystems. They are the main food of the dugong (sea cow), turtle and other marine animals and provide necessary surface for attachment, growth and development of many epiphytic seaweeds and small organism which in turn important food for many marine animals. Thus, sea grass as the “back bone” of complex marine communities but their role has often overlooked due to the submerged state.

Sea grass beds have long been viewed as unique and important coastal ecosystems. Much of the justification for the perceived importance of sea grass beds lies with its function as a nurturing habitat for fauna. Dense shelter in the form of sea grass canopy and food chains fueled by high in situ primary production, are held to provide the basis for very high faunal productivity (Larkum *et al.*, 1992).

In Peninsular Malaysia most of the sites of sea grass and coral ecosystem are situated in the East Coast. Sea grass areas are well distributed around Pulau Sibul and Pulau Besar, Mersing. There are three dominant species; that is *Enhalus acoroides*,

*Cymodocea serrulata* and *Halodule uninervis* (Rajudin, 1992). Other known site is in Besut, Terengganu and rarely occurred in Port Dickson.

Malaysia has been known among the most beautiful sites of coral reef. These ecosystems are located mostly in small island of the east coast, Sabah and Sarawak. In the west coast it can be found around Malacca Straits at Pulau Besar, Pulau Payar and Pulau Sembilan. Scuba diving, snorkeling and touring is the main activities in Pulau Redang, Pulau Tioman, Pulau Kapas and others due to the occurrence of their beautiful corals. Several areas such as Pulau Redang, Pulau Sibul, Pulau Besar, and Pulau Payar have been gazette as a Fisheries Protected Area and subsequently as a marine park. It is to provide the unique sea grass and coral reefs ecosystem unexploited marine gardens.

Most of popular tourist islands are found in east coast region and visits to these islands are usually made during the dry non-monsoonal months from March to October. The main Islands are Pulau Perhentian, Pulau Redang, Pulau Kapas, Pulau Aur, Pulau Pemanggil, and Pulau Tioman

## **2.2 Classification of Calanoid Copepods**

Sars first suggested the classification adopted for copepods order in 1900 to 1903 (cited in Zheng, 1989) who divided the Order Copepoda into seven suborders, Calanoida, Cyclopoida, Arguloida, Notodelphyoida, Monstrilloida and Cagilloida. Later a modified Sars's system of classification proposed by Gurney (1931-1933) is adopted, the raising of all the suborders to the rank of orders. To date, according to